




Contents lists available at ScienceDirect

Annals of Pure and Applied Logic

www.elsevier.com/locate/apal



The Kierstead's Conjecture and limitwise monotonic functions

Guohua Wu^{a,*}, Maxim Zubkov^{a,b,2}^a Division of Mathematical Sciences, School of Physical and Mathematical Sciences, Nanyang Technological University, 21 Nanyang Link, Singapore 637371, Singapore^b N.I. Lobachevsky Institute of Mathematics and Mechanics, Kazan Federal University, Kazan, Kremlevskaya 18, 420008, Russia

ARTICLE INFO

Article history:

Received 4 February 2016

Received in revised form 2 November 2017

Accepted 8 January 2018

Available online 11 January 2018

MSC:

03D45

03C57

Keywords:

Linear order

Limitwise monotonic function

Automorphism

ABSTRACT

In this paper, we prove Kierstead's conjecture for linear orders whose order types are $\sum_{q \in \mathbb{Q}} F(q)$, where F is an extended $0'$ -limitwise monotonic function, i.e. F can take value ζ . Linear orders in our consideration can have finite and infinite blocks simultaneously, and in this sense our result subsumes a recent result of C. Harris, K. Lee and S.B. Cooper, where only those linear orders with finite blocks are considered. Our result also covers one case of R. Downey and M. Moses' work, i.e. $\zeta \cdot \eta$. It covers some instances not being considered in both previous works mentioned above, such as $m \cdot \eta + \zeta \cdot \eta + n \cdot \eta$, for example, where $m, n > 0$.

© 2018 Elsevier B.V. All rights reserved.

1. Introduction

In this paper, we make progress toward solving a long-standing open problem of characterizing the order types of Π_1^0 -rigid computable linear orders (a linear order is Π_1^0 -rigid if it does not have any nontrivial Π_1^0 -automorphism). Downey's survey paper [1] provides an extensive motivation and research toward this problem. As usual, ω , ζ , η are the order types of natural, integers and rational numbers, respectively. Moreover, \mathbb{N} , \mathbb{Q} are the set of natural numbers and the set of rational numbers, correspondingly. By technical reasons, we assume that \mathbb{N} does not contain 0.

* Corresponding author.

E-mail addresses: guohua@ntu.edu.sg (G. Wu), maxim.zubkov@kpfu.ru (M. Zubkov).

¹ Wu is partially supported by MOE2011-T2-1-071 (M45110030), MOE2016-T2-1-083 (M4020333) from Ministry of Education of Singapore, and by AcRF grants RG29/14, M4011277 and RG32/16, M4011672 from Nanyang Technological University.² Zubkov is partially supported by the subsidy allocated to Kazan Federal University for the state assignment in the sphere of scientific activities, project No. 1.1515.2017/4.6, by RFBR grant 18-01-00574, 15-01-08252, and by MOE2011-T2-1-071 (M45110030) from Ministry of Education of Singapore, AcRF grants RG29/14, M4011277 from Nanyang Technological University.